

Diagnosis, and Management of Covid-19 Stage or Severity

	Asymptomatic or Presymptomatic	Mild Illness	Moderate Illness	Severe Illness	Critical Illness
Features	Positive SARS-CoV-2 test; no symptoms	Mild symptoms (e.g., fever, cough, or change in taste or smell); no dyspnea	Clinical or radiographic evidence of lower respiratory tract disease; oxygen saturation $\geq 94\%$	Oxygen saturation $< 94\%$; respiratory rate ≥ 30 breaths/min; lung infiltrates $> 50\%$	Respiratory failure, shock, and multiorgan dysfunction or failure
Testing	Screening testing; if patient has known exposure, diagnostic testing	Diagnostic testing	Diagnostic testing	Diagnostic testing	Diagnostic testing
Isolation	Yes	Yes	Yes	Yes	Yes
Proposed Disease Pathogenesis	<p>Viral replication (blue arrow) spans from Asymptomatic/Presymptomatic to Severe Illness. Inflammation (red arrow) spans from Moderate Illness to Critical Illness.</p>				
Potential Treatment	Antiviral therapy			Antibody therapy	
		Antibody therapy		Antiinflammatory therapy	
Management Considerations	Monitoring for symptoms	Clinical monitoring and supportive care	Clinical monitoring; if patient is hospitalized and at high risk for deterioration, possibly remdesivir	Hospitalization, oxygen therapy, and specific therapy (remdesivir, dexamethasone)	Critical care and specific therapy (dexamethasone, possibly remdesivir)